

2015

SCIENCE First Hand, V. 3(42)

The Epoch of Acceleration



Almost half a century ago, the prominent Russian physicist Gersh I. Budker, the founder of the Institute of Nuclear Physics (Novosibirsk), who pioneered and implemented the idea of increasing the energy of particle interactions by the colliding-beam method, called the charged particle accelerators the microscopes of modern physics. Indeed, these devices allow scientists to infer the structure of an object being observed from the way a particle flux is scattered on the object, the difference being that the scattered particles are not the quanta of light as in a microscope, but high-energy particles. The increase in the accelerators' limiting energy has revolutionized the theory of elementary particles and intranuclear interactions, and one can hardly overestimate the importance of this invention for science and technology.

At first, synchrotron radiation, which is emitted by charged particles moving with relativistic velocities in the magnetic field of accelerators, seemed an unfortunate by-product of the accelerating process because it meant large energy losses that had to be compensated. Since this radiation first occurred

in synchrotron, i.e., a cyclic resonance accelerator, it was called synchrotron radiation (SR), although its source can be any device that bends charged particles.

However, SR has surprisingly quickly made its way "from Cinderella to a princess" thanks to such "character traits" as a large width of the emission spectrum from infrared to hard X-rays, high directivity and polarization, periodicity on a nanosecond scale, and, finally, high power (although the latter traits are the merit of the special devices that generate SR). The idea of using the radiation of relativistic particles was proposed by the Russian theoretical physicist and future Nobel Prize winner Vitaly Ginzburg in 1947. Over the next half a century, SR became a very versatile and effective instrument for exploring the world.

There are three generations of SR sources. The first are the synchrotrons and storage rings designed for high energy physics. The second-generation sources are the accumulator rings that were designed specifically as SR sources. These sources generate SR with the help of bending magnets; since the radiation is directed tangentially to the particle path, like the light of car headlights when the car makes a turn, the beam has the shape of a fan with a large scattering angle.






Participants of the roundtable “New Horizons of Accelerator Physics: Present and Future of Bright Synchrotron Radiation Sources”: Prof. A. Seryi (John Adams Institute), Profs. A. Harrison and R. Walker (Diamond Light Source), Academician G. N. Kulipanov (BINP SB RAS), and the British Consul General J. Sharp.
Photo by the SIN–Russia, British Embassy in Russia

Judging by ancient chronicles, the mankind first encountered synchrotron radiation in 1054, when a bright “star”—the Crab Nebula, the result of a supernova explosion—lit up in the sky in broad daylight. The “inventor” of SR is the French physicist Alfred-Marie Liénard, who was the first to show, at the end of the 19th century, that an electron moving in a circular path intensively radiates electromagnetic waves. The first manmade synchrotron radiation was obtained in 1947 at the synchrotron of the company General Electric (United States); a few years later, synchrotron radiation was obtained at the synchrotrons of the Lebedev Physical Institute, Russian Academy of Sciences (Moscow, Soviet Union)

The Crab Nebula.
Image Credit: NASA, ESA, J. Hester, A. Loll (ASU)



RUSSIA AND BRITAIN AT A ROUNDTABLE

The SIN–Russia, based in the British Embassy in Moscow, is a part of the UK Science and Innovation Network, which is supported by the British government. With offices in 29 countries around the globe, this network contributes to the development of research and knowledge exchange, creates opportunities for cooperation between leading scientists and science organizations at the international level, and provides access to the large-scale international science infrastructure.

One of the SIN Russia’s long term goals is for Russia to be more strongly aware of the UK’s offer as a world-class place to carry out R&D in space science, composites, accelerator science, and arctic/polar research and innovation. Also, the SIN Russia works to encourage Russia to publish more of its research in international peer reviewed journals.

The SIN Russia organizes UK–Russian roundtables, science cafés, and lectures by leading British scientists in the major Russian cities, and takes part in science festivals and workshops. In 2014, the SIN-Russia worked closely with the UK Royal Society of Chemistry (RSC) to facilitate the first UK–Russia Chemistry Week in Moscow. One of the key issues of the event was the publication of research papers in the leading journals and open access to research publications. In 2014–2015, the SIN-Russia supported two UK–Russia science roundtables: Antarctic and Arctic in Focus of Science Research (together with Arctic and Antarctic Research Institute (St. Petersburg)) and New Horizons of Accelerator Technology. In 2015, we plan to organize events devoted to the problem of antimicrobial resistance, which is one of the priority issues for the UK Government.

Science and Innovation Network (SIN)–Russia, British Embassy in Russia


The third-generation SR sources include storage rings with long straight intervals and integrated magnetic structures of alternating polarity that generate SR with a smaller beam size, higher intensity, and much higher spectral brightness. The last parameter is the most important because it determines the magnitude of the useful photon flux. Thanks to the targeted effort of accelerator

physicists, the brightness of X-ray SR sources has been growing by three orders of magnitude every decade! Nevertheless, even in the most modern SR sources, the quantity of “useful” photons is only one thousandth of the full light flux; therefore, over the last decade, the world physical community has been working intensively to design SR sources of the next, fourth generation.

Roundtable participants Profs. A. Harrison and R. Walker (Diamond Light Source) and T. Yakovleva (SIN–Russia, British Embassy in Russia) speaking to editors of *SCIENCE First Hand*. Novosibirsk, 2015



Excursion to the INP World: Nine Years After



More than nine years ago, the staff of SCIENCE First Hand visited the Budker Institute of Nuclear Physics (INP), one of the largest national academic institutions and the internationally recognized science center. The fledgling editors of the newly organized journal, who were far from the world of high-level physics, were greatly impressed by seemingly endless labyrinths of accelerators underground, by amazing tangles of wires and obscure constructions, and by historical “round tables” established by Gersh I. Budker, the first director of the Institute.

Anyone interested could easily approach the Institute administration and attend the meetings of scientific councils. And of course, we could not but remember an amazingly great number of intelligent, interesting, and memorable men’s faces (sorry for chauvinism). Now, having got familiar with physical terms, we can say that here, in a relatively small area, it became possible to achieve a strikingly high concentration of high-energy intellectual particles. These impressions resulted in a series of reports about excursions to the INP world, where young physicists, postgraduates, and personnel of the Institute were our guides.

...Almost a decade later, we saw the same shiny old-fashioned coffee pots on the same round tables, the same hospitality, and many familiar faces of all different ages, who practically had not aged: being engaged in high-level physics seems to be beneficial not only to intellectual longevity.

However, tangles of wires have grown larger in already familiar underground “caves”, while the fully equipped experimental bunker for SR studies on the “ancient” VEPP-3 storage ring suggests stable associations with Herbert Wells’ underground world of Morlocks... Stability is appreciated when we consider traditions, continuity in research, and scientific creativity; however, on our next excursion to the national leader in high-energy physics, we hope to see the implementation of long desired plans for the radical reconstruction of the accelerating complex




2015

SCIENCE First Hand, V. 3(42)

© K. V. Zolotarev, P. A. Piminov,
A. D. Nikolenko, 2015

Pavel A. PIMINOV, Candidate of Physics and Mathematics, Director of the VEPP-3/VEPP-4 storage ring system at the Budker Institute of Nuclear Physics, Siberian Branch, Russian Academy of Sciences (Novosibirsk); author and coauthor of 103 scientific publications

Konstantin V. ZOLOTAREV, Candidate of Physics and Mathematics, Head of Division at the Budker Institute of Nuclear Physics, Siberian Branch, Russian Academy of Sciences (Novosibirsk); Associate Professor at the Accelerator Physics Division, Novosibirsk State University; author and coauthor of 93 scientific publications



SR at the INP: formula of success

It is known that synchrotron radiation (SR) is the daily bread for many users; for accelerator physicists, it is a parasite because it takes away a large part of energy pumped with great effort into the beam of charged particles accelerated nearly to the speed of light. Besides, it is hard work, though the end results are sometimes less than the efforts spent to achieve them. The point is that now the SR sources at the Institute of Nuclear Physics (INP) are VEPP-3 (created in 1972) and VEPP-4M (put in operation in the early 1980s and later modernized), on which studies in the field of elementary particle physics are conducted. Consequently, SR experiments take only 15% of the total time of accelerator operation.


Thus, although SR experiments have been carried out at the INP ever since 1973, they still use VEPP-3/VEPP-4, which are not sufficiently bright first-generation SR sources operating in the X-ray range (the wavelength varying from 0.01 to 1 nm) and a beam energy of 2 or 4 GeV.

It is worth mentioning that the 2003 commissioning of the first line of a free electron laser system (a source of powerful terahertz radiation beams) principally expanded the INP's research potential; however, it is still necessary to develop a more powerful new generation SR source that will operate in the X-ray range.

On December 1, 1981, the Siberian Center for Synchrotron Radiation was organized in order to coordinate the efforts directed to developing SR studies, using SR sources efficiently, and improving the quality of research. The center was based on the acceleration equipment and laboratories of the Budker Institute of Nuclear Physics SB RAS. In 1991, it was reorganized into the Siberian International Center for Synchrotron Radiation, an open laboratory for Russian and foreign organizations and individuals. In 2003, the first line of a free electron laser was put in operation. In 2005, the Center was renamed the Siberian Center for Synchrotron and Terahertz Radiation

At the INP, VEPP-3/VEPP-4 are used in the acceleration storage complex for SR generation, VEPP-3 being a booster (intermediate) accelerator for the VEPP-4 collider (diagram on the right). Acceleration occurs in the range of energy from 360 MeV to 2 GeV; in the storage mode the accelerator can hold beams with an energy of 2 GeV and a current of about 100 mA for a long time (5–6 h). It is in this mode that the work using SR is conducted.

Upper right: the rectilinear part of the VEPP-3 storage ring; *lower right:* the experimental chamber for SR on the VEPP-4 storage ring





Today, several tens of large research centers with third-generation SR sources, such as Diamond Light Source (United Kingdom), Swiss Light Source (Switzerland), Soleil (France) and others are in operation; yet another two “brighter” sources—NSLS-2 (United States) and MAX-IV (Sweden)—are being commissioned. All the centers are in high demand and work at maximum capacity; there are expert systems to allocate the “beam time”: on a competitive basis, for researchers, and for a fee, for other users. Typical fee-paying users are large pharmaceutical companies that invent and test new drugs, which is hard and costly work that never ends.

The INP cannot compete with the large centers on these “highways” because of the beam characteristics of the available sources. So, if our scientists or technologists have a particular problem, e.g. testing a new drug, it is easier for them to go to England or to France, to the European Synchrotron Radiation Facility (ESRF), in which our country participates.

Nonetheless, our sources have been heavily exploited for both research and routine technological aims. For instance, the researchers from the Institute of Catalysis SB RAS (Novosibirsk) are constantly analyzing samples of novel catalysts intended for industrial applications. Our main advantage is that SR at the INP has largely maintained its initially unrestricted status of a research tool; so, practically any scientist can use it to verify his/her idea (however crazy it may be).

In this respect, it is very important that our SR sources are located in such an unusual infrastructure as Novosibirsk Akademgorodok, i.e. in a vast multidisciplinary environment. For instance, archaeologists, who work in the same neighborhood, can ask physicists to analyze any artifact they are interested in. Indeed, new knowledge is gained, as a rule, by combining unique samples

In the bunker for SR from the VEPP-3 storage ring, the work is organized as in a famous Russian fairy tale: the more, the merrier. In an experimental hall there are ten user stations located in an area that is only 90 m²; so, the room is overflowing with equipment. *Upper right:* the unique first station Detonation, which consists of an explosion chamber and a detecting unit, where it is possible to operate with explosives weighing up to 50 g

This is the living history of accelerator physics: the B-4 booster synchrotron in which there occurs initial acceleration of electron (positron) beams to the injection energy (360 MeV) into the VEPP-3 storage ring. It is likely to be the only “antique” working synchrotron in the world



At present, the Siberian Center for Synchrotron and Terahertz Radiation has 12 SR stations and 4 terahertz radiation stations. The major aims and tasks of the center are fundamental and applied studies in physics, chemistry (including catalysis), biology, medicine, ecology, geology, materials science, and also the development of new methods and technologies as well as the creation of new specialized radiation sources and experimental stations

provided by users and adequate research tools which can be realized with the use of SR.

Moreover, we elaborate the procedures which in principle are difficult to develop at large SR centers, which, to some extent, is due to administrative restrictions. A good example is the study of detonation processes with a submillisecond time resolution in a special explosion chamber located at the SR output channel. Since SR has the form of short flashes resembling the time structure of short electron bunches rather than a continuous flux (in our case, the duration of these flashes is 1 ns, and the recurrence period is about 100 ns), by studying the features of the interaction between radiation and the substance we can determine the current state of the substance with a corresponding time resolution. It means that during detonation it is possible to examine the character of chemical processes occurring in the moving detonation front, the growth dynamics of detonation nanodiamonds, and other effects that interest the specialists.

The first experimental station, named Detonation, was set on the VEPP-3 storage ring, and later a second station started its work on the VEPP-4 storage ring: the new chamber made it possible to investigate the detonation of charges weighing up to 200 g. Now this station is being modernized in order to study the effect of powerful laser plasma pulses on construction materials. The knowledge of these processes will be necessary for designing future thermonuclear reactors.

Other user stations are regularly updated as well. Thus, the application of new focusing lenses provided the improvement of the spatial resolution of X-ray fluorescent elemental analysis, which helps to determine not only the chemical composition of a sample but also the spatial arrangement of separate elements. Although we cannot compete with other SR centers in this field, we have succeeded in obtaining some interesting results. For instance, researchers from the Limnological Institute SB RAS (Irkutsk) detected a response of the elemental composition of bottom sediments to a change in the climate,

such as Milankovich cycles, which allows researchers to use SR to investigate paleoclimate. Similar work is in progress now with bottom sediments from Altai lakes, in particular, Lake Teletskoye.

Yet another method of studying the state of matter under extreme conditions (at ultrahigh pressures of several GPa and temperatures up to 1000°C) is the diamond anvil technique, in which a sample is squeezed between two diamond nibs. Thereby it is possible to reach ultrahigh pressures in a small volume, thus modeling the behavior of matter at large depths, i.e. in the Earth's mantle or even at the center of the planet. This station belongs to the Institute of Solid State Chemistry SB RAS (Novosibirsk).

Our latest developments include the COSMOS metrological station, designed for testing satellite devices, and PLAMYA, planned for studying rapid reactions such as combustion. The latter is being created in collaboration with the Institute of Chemical Kinetics and Combustion SB RAS (Novosibirsk).

However, the INP in the SR world is not just a participant: the Institute is also an active developer. The INP is practically a monopolist in producing superconducting wigglers (multipolar magnets generating a sign-variable periodic magnetic field), which are installed in rectilinear spaces of electron storage rings to increase radiation intensity. Novosibirsk physicists and engineers control the whole production cycle of these complex devices: from designing and manufacturing to testing and assembly on site. All over the world, from Australia to Brazil and North America, more than 20 wigglers made in Novosibirsk are currently in use. The INP researchers have developed, produced, and installed superconducting devices practically in all world SR centers, including Spring-8 (Japan), ELETTRA (Italy), CLS (Canada), synchrotrons in Brazil and Australia, and the only Russian specialized SR source at the Kurchatov Institute in Moscow.


At present, the research group dealing with wigglers is engaged in producing undulators, i.e. superconducting devices with a large number of poles and a low magnetic field. In contrast to wigglers, in undulators the radiation from separate poles is coherent, which provides monochromatic radiation with a substantially higher spectral brightness. All modern centers are interested in these devices. For instance, there is a tentative agreement on collaboration in this field with DLS (United Kingdom).

The main problem of the Siberian Center for Synchrotron and Terahertz Radiation has been the absence of its own specialized SR source, although at least five different ways to create it have been proposed in the last decade. The INP has the necessary experience, technologies, and production facilities; the only thing missing is financing.



New station for studying detonation processes, planned for explosives weighing up to 200 g, in the VEPP-4 SR hall

Radiation box of the X-Ray Phase Contrast Microscopy and Microtomography station operating with SR beams from the VEPP-4 storage ring; at present, it is used for experiments on microbeam therapy of carcinomas. Specialists from the Institute of Cytology and Genetics SB RAS (Novosibirsk) are also participating in this work. At the current stage, the experiments are carried out with laboratory animals



This superconducting wiggler with a magnetic field of up to 3 T and a cost of about 1.4 million euro was manufactured at the INP for the ANKASR source (Germany). The work took about one and a half years. This device, for the first time, used a cryostat with indirect cooling of the magnetic system that provided the safety, efficiency, and simplicity of the cryogen service. It is planned to use similar wigglers in the CLIC linear collider in CERN

The latest version of the new source differs from all the previous (and rejected) ones in that it is the most cost-effective. The project plans to use the tunnel in which VEPP-3 is currently located. It is also planned to extend the existing experimental chamber and to place new user stations there. A superconducting wiggler and a pair of superconducting dipole magnets will be used as radiation sources. A special magnetic structure of the ring will combine extreme compactness with a possibility for optimizing the beam brightness.

In conclusion, it is worth noting that in the last decades interest in the studies at the interface of sciences has greatly increased worldwide. In our academic center, a multidisciplinary scientific community that unites researchers from the institutes of Novosibirsk

Akademgorodok and from other science centers has formed. The sincere interest of these specialists in SR studies guarantees a flood of research materials and publications, as well as extremely efficient use of all available methods and instruments. Novosibirsk National Research University must also be interested in a new SR source: our SR Center can train the students of all natural science departments, as is done at many other universities worldwide.

The INP has long deserved the right not to be a shoemaker's child without shoes: it must have its own specialized SR source at the Siberian Center for Synchrotron and Terahertz Radiation. The only things lacking are federal financing and political will. Against all odds, we remain optimistic and believe in the future.